



Research papers

The role of cave ventilation in the triple oxygen and hydrogen isotope composition of condensation waters in Altamira Cave, northern Spain

Fernando Gázquez^{a,b,*}, Luis Quindós^{c,d}, Daniel Rábago^c, Ismael Fuente^c, Santiago Celaya^c, Carlos Sainz^{c,d}

^a Department of Biology and Geology, Universidad de Almería, Carretera de Sacramento s.n, La Cañada de San Urbano, Almería 04120, Spain

^b Andalusian Centre for the Monitoring and Assessment of Global Change (CAESCG), University of Almería, Spain

^c Radon Research Group, Faculty of Medicine, University of Cantabria, Avda. Cardenal Herrera Oria s/n, E-39011 Santander, Spain

^d The Cantabrian International Institute for Prehistoric Research (IIIPC), Spain

ARTICLE INFO

This manuscript was handled by Marco Borgia, Editor-in-Chief, with the assistance of Huade Guan, Associate Editor

Keywords:

Cave management
Cave monitoring
Condensation water
Stable isotopes
Oxygen-17
Triple oxygen isotopes

ABSTRACT

In cave environments, water vapor condensation occurs naturally when warmer/wet air masses flow close to colder cave surfaces. Artificial microclimate perturbations in show caves can enhance this process, leading to potential deterioration of rock art and degradation of speleothems. Here we investigate the triple oxygen and hydrogen isotopic compositions of condensation water in Altamira Cave (Cantabria, northern Spain) to evaluate the potential of stable isotopes in the study of condensation mechanisms in caves. We assess the role of cave ventilation in the spatiotemporal isotopic variability of condensation water in Altamira Cave. To this end, water drops that condense naturally on artificial supports in different parts of the cave were collected for 7 years and their isotopic compositions ($\delta^{17}\text{O}$, $\delta^{18}\text{O}$, δD and derived parameters ^{17}O -excess and d-excess) were compared to those of droplets with no apparent dripping taken from the cave ceiling (i.e. presumably condensation water) and fast dripping points (i.e. infiltration water) during the same period. Condensation waters in the outmost cave sectors, closer to the entrance, show higher $\delta^{17}\text{O}$, $\delta^{18}\text{O}$ and δD values during the cave ventilation period (June to October) compared to the rest of the year. This seasonal pattern can be explained by changes in the contributions of two moisture sources for condensation: advection of allochthonous water vapor from outside during the cave ventilation period and recycling of autochthonous vapor generated from cave dripwater during the stagnation period. In contrast, the isotopic values of condensation waters in the inner cave sectors are similar to those of infiltration water, with insignificant seasonal variability. This suggests that water condensation in the inner cave sectors is sourced by autochthonous vapor, with no significant contributions of external moisture, even during the cave ventilation period. We conclude that allochthonous water vapor condenses preferentially in the *Entrance Hall* and does not affect significantly the rest of the cave. These results are relevant for the management of Altamira Cave and for future investigations on condensation mechanisms in cavities elsewhere.

1. Introduction

The prehistoric paintings of the *Polychrome Hall* of Altamira Cave are the maximum expression of the Paleolithic rock art in the Iberian Peninsula. This cave hosts priceless rock art paintings, some 36,160 – 15,329 cal. BP that have earned it the name “Sistine Chapel” of the Paleolithic, due to the reddish and ochre-colored bison and deer paintings that cover the ceiling of the renowned *Polychrome Hall* (Pike et al., 2012; Gázquez et al., 2017a) (Fig. 1). The cave was declared World

Heritage Site by the UNESCO in 1985 (<http://whc.unesco.org/en/list/310>) and was recently reopened to the public under an extremely restrictive visitor regime (Gázquez et al., 2016, 2017a; Sainz et al., 2017). Preserving the integrity of these and others rock art features from natural and anthropic perturbations is a priority in the management of show caves.

Among the several threats to the rock art heritage, resuspension and mobilization of pigments (mostly ochres and charcoal) by cave water, including seepage and condensation waters, are major hazards. Water

* Corresponding author at: Department of Biology and Geology, Universidad de Almería, Carretera de Sacramento s.n, La Cañada de San Urbano, Almería 04120, Spain.

E-mail address: f.gazquez@ual.es (F. Gázquez).

<https://doi.org/10.1016/j.jhydrol.2021.127416>

Received 4 May 2021; Received in revised form 20 November 2021; Accepted 28 December 2021

Available online 7 January 2022

0022-1694/© 2022 Elsevier B.V. All rights reserved.

films and drops can accumulate in some parts of the cave walls and ceilings. Identifying the source of water (condensation vs. infiltration) is crucial to develop protocols devoted to minimize the impact on the cave features, including rock art and speleothems (Dublyansky and Dublyansky, 2000; Fernández-Cortés et al., 2006).

The largest intakes of external water vapor in Altamira Cave may occur during the cave ventilation period (June to October) (Cuezva et al., 2009; García-Antón et al., 2013; Quindós et al., 1987; Gázquez et al., 2016; Sainz et al., 2017), when relatively warm and moist air masses enter the cave and cool, forcing condensation. In order to reduce the inputs of airborne particles (i.e. organic matter, fungi and bacteria) that could deteriorate the Paleolithic rock art of Altamira Cave, an artificial metal door was installed in 2008 between the *Entrance Hall* (~20 m apart from the entrance) and the rest of the cave. This intervention has had a positive impact on the microclimate stability of the inner cave sectors in terms of temperature stability and decrease in airborne particles (Sainz et al., 2018); however, the role of this artificial barrier in the condensation processes in the cave has not been investigated to date. Indeed, specific studies on water condensation in Altamira Cave have not been performed yet.

Direct measurements of the condensation/evaporation processes have been conducted previously in caves worldwide by using suspended glass plates (Sarbu and Lascu, 1997), lysimeters and metallic devices (Dublyansky and Dublyansky, 1998; Gázquez et al., 2015a; 2017b) and refrigerated containers (Tarihule-Lips and Ford, 1998; Liñán et al., 2021), in order to quantify the magnitude of these mechanisms. Electrical devices coupled to data loggers have been utilized also to quantify condensation in caves (De Freitas and Schmekal, 2006). In addition, indirect measurements of the effects of condensation have been

conducted by monitoring weight loss of rock tablets (Calaforra, 1996; Klimchouk, 1996; Tarihule-Lips and Ford, 1998; Klimchouk and Aksem, 2002; White et al., 2021), using micro-erosion meter measurements of cave surface retreatment (Calaforra et al., 1993; Klimchouk, 1996; Gázquez et al., 2015a) and determining thickness and age of weathering rinds (Auler and Smart, 2004). Indirect estimates of condensation/evaporation can be also obtained from microclimate monitoring of cave atmosphere and rock temperatures (De Freitas and Schmekal, 2003; Fernández-Cortés et al., 2006). Furthermore, theoretical approaches to the condensation mechanisms in caves have been developed in the last decades (Dreybrodt et al., 2005; Gabrovšek et al., 2010). More recently, thermal image analyses have been used to quantify condensation-evaporation processes in caves (Liñán et al., 2021).

The oxygen and hydrogen isotopes composition of water [$^{18}\text{O}/^{16}\text{O}$ and $^2\text{H}/^1\text{H}$ or $\delta^{18}\text{O}$ and δD , respectively, when standardized to the international water standard Vienna-Standard Mean Ocean Water (V-SMOW)] has been widely used to track a variety of processes in the hydrosphere (Bowen et al., 2019 and references therein), including recent preliminary studies of condensation water in caves by Liñán et al. (2021). These authors found that condensation waters in Nerja Cave, southern Spain, display higher $\delta^{18}\text{O}$ and δD values and lower d-excess values during the cave ventilation periods than during the rest of the year. This seasonal pattern was attributed to the impact of evaporation on condensation water, which takes place during the cave ventilation period because of low relative humidity in the cave atmosphere (60–80%).

In addition to the traditional measurements of $\delta^{18}\text{O}$ and δD in waters, recent analytical advances have permitted obtaining precise measurements of the triple oxygen isotope composition ($\delta^{17}\text{O}$ and $\delta^{18}\text{O}$) (Barkan




Fig. 1. A. Location of Altamira Cave and distribution of the sampling sites in the cavity, including artificial supports to collect condensation water (C1, C2 and C3); B. View of the ceiling of Polychrome Hall (location is marked in orange rectangles in panels 1A and 1C) with collection locations for the dripwater (D1) and ceiling droplets (D2-D5) labelled; C. Vertical profile of Altamira Cave and location of the sampling sites (modified from García-Antón et al., 2014).

and Luz, 2005; Steig et al., 2014). The $\delta^{17}\text{O}$ deviations with respect to the $\delta^{17}\text{O}$ – $\delta^{18}\text{O}$ Global Meteoric Water Line (GMWL), which has a proposed slope of 0.528, are expressed as ^{17}O -excess [^{17}O -excess = $\ln(\delta^{17}\text{O}/1000 + 1) - 0.528 \ln(\delta^{18}\text{O}/1000 + 1)$] (Luz and Barkan, 2010). This reference slope is commonly used to express triple oxygen deviation in the hydrological cycle (Aron et al., 2021; Surma et al., 2021).

The ^{17}O -excess parameter in rainfall has been found to be sensitive to variations in relative humidity at the moisture source, with smaller temperature dependence than d-excess ($\text{d-excess} = \delta\text{D} - 8 \cdot \delta^{18}\text{O}$) (Barkan and Luz, 2005, 2007; Luz and Barkan, 2010; Uechi and Uemura, 2019). Likewise, other parameters, including raindrop re-evaporation and moisture recycling during water vapor transport can modulates the final ^{17}O -excess signal of rainwater (Tian et al., 2018; Giménez et al., 2021). Thus, measuring the triple oxygen and hydrogen isotope composition of water can help to track environmental conditions (e.g. temperature vs. relative humidity) during water phase changes and during transport of water vapor masses (Aron et al., 2021; Surma et al., 2021).

Here, we conducted a 7-years monitoring study of stable isotopes ($\delta^{17}\text{O}$, $\delta^{18}\text{O}$, $\delta^2\text{H}$ and derived parameters d-excess and ^{17}O -excess) in condensation water and dripwater in Altamira Cave. We aim to identify isotopic differences between seepage and condensation water, in order to assess whether the numerous droplets observed in the ceiling of the *Polychrome Hall* of Altamira Cave are associated with either condensation or infiltration water. In addition, we investigate the origin of vapor for water condensation and the spatiotemporal pattern of the condensation mechanisms in this cave. To this end, we analyzed waters deposited on artificial supports in several locations of the cavity, as well as dripwater from fast drips (i.e. discharge point related to seepage of meteoric water) and ceiling droplets with no apparent dripping (presumably associated with water condensation) in the *Polychrome Hall*. The microclimate characteristics related to cave atmosphere ventilation (air temperature, relative humidity and CO_2 content) were monitored during the length of the study, in order to characterize the parameters that control the cave ventilation pattern and to identify connections with the condensation processes. We use an isotopic mixing model to demonstrate that different sources of water vapor contribute to condensation water in Altamira Cave and that the relevance of these sources change temporally and spatially.

2. Setting and cave description

Altamira Cave ($43^\circ 22' 37''\text{N}$; $4^\circ 07' 11''\text{W}$) is located near the village of Santillana del Mar (Cantabria, northern Spain), at an elevation of 152 m a.s.l and 4.5 km from the Cantabrian Sea. The cavern runs NW-SE and has developed along a subhorizontal sequence of Cenomanian-Turonian marine carbonates (Hoyos et al., 1981) where the difference in level between the entrance and its deepest part (*Grave Hall*) is barely 26 m. Several cave sections can be distinguished according to their morphology and distance/depth from the entrance: the *Entrance Hall* is a ~20 m-long horizontal hall; from there, access to the rest of the cave is via an artificial door located in a narrow passage, ~2 m wide. Beyond, the cave divides along two branches. To the east is the *Polychrome Hall*, ~15 m long, ~7 m wide and up to 3 m high and lying about 2 m lower than the cave entrance. The other branch gives access to the *Walls Hall*, roughly 50 m in length, 20 m wide and up to 7 m high; located ~6 m below the entrance.

A passage, roughly 60 m long runs SE-NE into the *Great Hall*. From here, artificial stairs carved in the rock lead to the *Grave Hall*, lying ~15 m below the cave entrance and containing an ephemeral water pool that only appears during periods of high dripwater discharge. To the South, along a 10 m-long passage, lies the *Shower Hall*. This in turn leads into the *Well Hall*, at the depth of 12 m beneath the cave entrance. Finally, there is a meandriform sub-horizontal passage (called the *Horsetail*) approximately 40 m long that represents the innermost cave area. The thickness of the cap rock overlaying the cave varies from 2 to 18 m (Elez et al., 2013) (Fig. 1). There is no permanent running watercourse inside

the cave.

The climate in this region is Oceanic (Cfb climate type, according to the Köppen climate classification). The rainfall is generally abundant, exceeding 1,000 mm/year. July is the driest month (30 mm on average) and the rainiest period occurs from October to December (Fig. 2). The mean external air temperatures are minimum in January (9 °C) and maximum in July (21 °C). The vegetation coverage over the cave is dominated by grasses with minor presence of scrubs and trees (Fig. 1A).

Altamira Cave has a relatively stable microclimate, with air temperatures that vary <2 °C throughout the year in the innermost areas (*Well Hall* and *Horsetail*) and up to 5 °C in the outermost part of the cave (*Entrance Hall*) (Gázquez et al., 2016). The mean temperature in the *Polychrome Hall* during the period of study was 14.2 ± 0.5 °C (maximum value of 15 °C and minimum value of 13.2 °C).

The air temperature and ventilation regime of Altamira Cave are fundamentally controlled by the annual thermal oscillations of the outside air temperature and the water saturation of the epikarst and the soil over the cave (Cuezva et al., 2011). The outside temperature oscillations are transmitted into the cave through the rock, which attenuates its amplitude and causes a lag between the outer thermal wave and the temperature of the subterranean atmosphere (Elez et al., 2013). Consequently, the cave ventilation period occurs in summer (June to October), while the cave stagnation period takes place in winter (November to May). The annual variation of in-cave air temperature exhibits a sinusoidal pattern with a thermal lag time (~2 months on average) between the innermost and outermost cave areas that depends on the outside temperature and the thickness of the rock over the different cave sectors (Cuezva et al., 2011; Quindós et al., 1987; Sainz et al., 2018).

3. Methods

We assess the spatial gradient and temporal variability of the isotopic composition of condensation water and dripwater, as well as cave environmental parameters (air temperature, relative humidity and CO_2 content) in Altamira Cave, by monitoring and repeatedly sampling from the cave entrance to more stable locations deeper within the cave.

3.1. Water sampling

From November 2014 to March 2017, condensation water that accumulates on a metallic door that separates the *Entrance Hall* and the rest of the cave (site C1 in Fig. 1) was collected using disposable pipets (~200 µl) (n = 11). From March 2017 to October 2020, sampling was conducted approximately every two weeks in the same location, but condensation was collected from the metal support of a thermometer that is 1 m apart from the door. In addition, condensation water from two inner cave locations (sites C2 and C3 in Fig. 1) was collected from metal supports of thermometers using the same method from March 2017 to August 2020. Water samples were stored at 4 °C in 2 ml glass tubes provided with a micro-insert (~300 µl; Thermo Fisher, n. 10376102) and capped with silicone septum caps. A total of 76 surveys were conducted and a total of 192 condensation water samples from artificial supports were collected.

Dripwater samples from a relatively fast discharge point (maximum of 143 ml/day; mean of 61 ± 21 ml/day) in the *Polychrome Hall* (site D1 in Fig. 1) were collected approximately every two weeks between March 2017 and August 2020. A graduated container was placed under the drip point and volumes were recorded between surveys in order to calculate drip rates. Water samples were collected from this container and stored in 14 ml glass tubes at 4 °C. A total of 76 dripwater samples were collected from site D1. In addition, water droplets from four selected locations of the ceiling of the *Polychrome Hall* (i.e., sites with no visibly dripping; D2-D5 in Fig. 1) were collected and stored using the same protocol as for condensation water. A total of 254 samples from these four locations were taken every two weeks between March 2017 and




Fig. 2. Temporal variations of oxygen and hydrogen stable isotopes (A. $\delta^{18}\text{O}$; B. δD ; C. d-excess; D. ^{17}O -excess) of condensation water and dripwater in Altamira Cave (site C1 in Fig. 1). Results of modeled isotope composition of condensation water at site C1 are displayed (dashed lines). Relative humidity (E) (site C1), CO_2 concentration in the cave atmosphere (F) (site C1), air temperature in the cave (site C1) and outside (G) and monthly rainfall in Santillana del Mar (H) are presented. A 2-point moving average has been used to represent air temperature. The yellow bands represent cave ventilation periods (June to October).

October 2020.

3.2. Stable isotopes analysis

Samples collected between July 2013 and July 2014 were analyzed for $\delta^{18}\text{O}$ and δD using a L1102-i Picarro water isotope analyzer at the Godwin Laboratory for Paleoclimate Research, University of Cambridge, UK. Samples collected between August 2014 and October 2020 were analyzed for $\delta^{17}\text{O}$, $\delta^{18}\text{O}$ and δD at the Laboratory of Stable Isotopes of the University of Almeria, Spain using a L2140-i Picarro water isotope analyzer (Picarro, Inc., Santa Clara, California, USA).

Each sample was injected ten times into the vaporizer (A0211 manufactured by Picarro, Inc.), which was heated to 110 °C. Potential contamination produced by organic compounds in water, which may affect specially the measurements of ^{17}O -excess values, was removed by using a Picarro micro-combustion module (MCM) coupled to the L2140-i Picarro analyzer (Gázquez et al., 2015b). Memory effects from previous samples were avoided by rejecting the first three analyses. Values for the final 7 injections were averaged with a typical mean instrumental precision ($\pm 1\text{SD}$) of $\pm 0.02\text{‰}$ for $\delta^{17}\text{O}$, $\pm 0.03\text{‰}$ for $\delta^{18}\text{O}$ and $\pm 0.19\text{‰}$ for δD when using the L2140-i Picarro analyzer, as observed from repeated analysis of an in-house water standard ($n = 54$) together with the water samples. The instrumental precision of the L1102-i Picarro analyzer was $\pm 0.05\text{‰}$ for $\delta^{18}\text{O}$ and $\pm 0.6\text{‰}$ for δD . The in-run drift of the instruments was monitored by analyzing a water standard (BOTTy) every 6 samples. The results were normalized against V-SMOW-SLAP in the case of triple oxygen isotopes, and against V-SMOW-SLAP-GISP in the case of hydrogen isotopes, by analyzing internal standards before and after each set of fifteen to twenty samples (see Supplementary Table 1 for examples of our calibration). To this end, three internal water standards (JRW, $\delta^{17}\text{O} = -9.99\text{‰}$, $\delta^{18}\text{O} = -18.90\text{‰}$ and $\delta\text{D} = -146.5\text{‰}$; BOTTy, $\delta^{17}\text{O} = -3.88\text{‰}$, $\delta^{18}\text{O} = -7.40\text{‰}$ and $\delta\text{D} = -50.38\text{‰}$; SPIT, $\delta^{17}\text{O} = -0.09\text{‰}$, $\delta^{18}\text{O} = -0.15\text{‰}$ and $\delta\text{D} = -0.44\text{‰}$) were calibrated previously for triple oxygen isotopes against V-SMOW and SLAP by a two-points calibration, using $\delta^{17}\text{O}$ of 0.0‰ and -29.69865‰ , respectively, and $\delta^{18}\text{O}$ of 0.0‰ and -55.5‰ , respectively (Schoenemann et al., 2013). This approach assumes V-SMOW and SLAP have ^{17}O -excess = 0. The δD of the internal standards was calibrated against V-SMOW, GISP and SLAP using a three-points calibration. Note that GISP does not have an accepted $\delta^{17}\text{O}$ value, so cannot be used for triple oxygen isotope calibration. Instead, the GISP water standard was analyzed as an unknown during the calibration of our internal standards, yielding an average ^{17}O -excess value of 19 ± 18 per meg. This value is in good agreement with the results reported by previous studies (e.g. Schoenemann et al. 2013; 22 ± 11 per meg).

The ^{17}O -excess was calculated for each injection using the normalized $\delta^{17}\text{O}$ and $\delta^{18}\text{O}$ values. The final reported value is the average of ^{17}O -excess values from all 7 suitable injections. All ^{17}O -excess values are given in per meg units (1 per meg = 0.001‰). Typical in-sample ^{17}O -excess and d-excess precisions (1SD) in water standards (BOTTy, $n = 54$), were 13 per meg and 0.2‰, respectively, similar to the precision obtained by other CRDS instruments (Wassenaar et al., 2021).

3.3. Microclimate monitoring

Measurements of air temperature and air CO_2 concentrations were conducted at sites C1, C2 and C3, in the Polychrome Hall (Fig. 1) and outside the cave close to the entrance door, between January 2014 and October 2020. Rainfall amount was recorded by a meteorological station located in the terrain over the cave during the length of the study. A total of 191 monitoring campaigns were conducted during this period, typically every two weeks and between 10 a.m. and 12 p.m. The in-cave temperature measurements used mercury thermometers with a precision of ± 0.05 °C that were installed in the cave at the beginning of this study. For the measurements of the external air we used a portable digital temperature probe (Vaisala HMP155) with a precision of ± 0.1 °C. The concentration of CO_2 in air was analyzed in situ using a Testo

Table 1

Summary of triple oxygen and hydrogen isotopes in waters from Altamira Cave (*see Fig. 1 for sampling site locations). Results of the cave ventilation period (V; June to October) and stagnation period (S; November to May) are averaged.

Sampling site	Location code*	Period	Type	V/ S	$\delta^{17}\text{O}$ (‰) \pm 1SD	$\delta^{17}\text{O}$ (‰) max	$\delta^{17}\text{O}$ (‰) min	$\delta^{18}\text{O}$ (‰) \pm 1SD	$\delta^{18}\text{O}$ (‰) maxO (‰) max	$\delta^{18}\text{O}$ (‰) min	δD (‰) \pm 1SD	δD (‰) max	δD (‰) min	d- excess (‰) \pm 1SD	d- excess (‰) max	d- excess (‰) min	^{17}O - excess (per meg) \pm 1SD	^{17}O - excess (per meg) max	^{17}O - excess (per meg) min
Entrance Hall	C1	15/07/ 2013 to 12/08/ 2020	Condensation on artificial support	V	-2.39 \pm 0.37	-1.39	-3.18	-4.60 \pm 0.69	-2.66	-6.05	-22.4 \pm 4.6	-13.9	-32.7	14.3 \pm 2.1	17.2	7.4	27 \pm 12	60	7
				S	-2.96 \pm 0.21	-2.96	-3.23	-5.65 \pm 0.40	-4.56	-6.22	-30.0 \pm 3.0	-20.6	-32.7	15.2 \pm 1.1	17.3	12.1	28 \pm 9	58	13
Middle Sector	C2	13/03/ 17 to 12/08/ 2020	Condensation on artificial support	V	-3.14 \pm 0.13	-2.87	-3.36	-5.99 \pm 0.25	-5.49	-6.38	-33.5 \pm 1.2	-31.9	-36.6	14.3 \pm 1.1	15.8	11.8	28 \pm 9	42	8
				S	-3.12 \pm 0.14	-2.90	-3.35	-5.95 \pm 0.26	-5.53	-6.39	-32.7 \pm 1.9	-29.9	-37.0	14.8 \pm 1.2	17.3	11.8	29 \pm 8	48	12
Great Hall	C3	13/03/ 17 to 12/08/ 2020	Condensation on artificial support	V	-3.14 \pm 0.13	-2.84	-3.40	-5.99 \pm 0.25	-5.42	-6.47	-33.3 \pm 1.1	-31.3	-37.1	14.6 \pm 1.3	17.6	12.0	26 \pm 10	47	11
				S	-3.13 \pm 0.16	-2.78	-3.44	-5.97 \pm 0.31	-5.29	-6.56	-32.8 \pm 2.0	-29.2	-37.3	14.9 \pm 1.2	16.9	12.8	30 \pm 11	54	12
Polychrome Hall	D1	13/03/ 17 to 12/08/ 2020	Dripwater	V	-3.08 \pm 0.08	-2.86	-3.18	-5.87 \pm 0.14	-5.48	-6.05	-32.8 \pm 0.4	-31.6	-33.6	14.1 \pm 0.8	14.9	11.9	22 \pm 7	36	11
				S	-3.10 \pm 0.04	-3.00	-3.16	-5.91 \pm 0.07	-5.71	-6.02	-32.8 \pm 0.4	-32.2	-33.7	14.5 \pm 0.4	15.0	12.8	26 \pm 7	39	11
Polychrome Hall	D2	22/03/ 17 to 12/08/ 2020	Droplets on the cave ceiling	V	-2.95 \pm 0.12	-2.73	-3.18	-5.63 \pm 0.23	-5.20	-6.02	-31.2 \pm 0.85	-29.4	-32.5	13.9 \pm 1.1	15.7	11.4	25 \pm 9	44	5
				S	-3.03 \pm 0.10	-2.83	-3.26	-5.78 \pm 0.19	-5.36	-6.22	-31.5 \pm 0.9	-29.6	-33.7	14.7 \pm 1.1	16.9	12.1	26 \pm 11	61	3
Polychrome Hall	D3	22/03/ 17 to 12/08/ 2020	Droplets on the cave ceiling	V	-2.96 \pm 0.14	-2.47	-3.12	-5.64 \pm 0.26	-4.71	-5.96	-31.5 \pm 0.8	-29.2	-33.0	13.7 \pm 1.4	15.8	8.3	25 \pm 8	51	14
				S	-3.00 \pm 0.13	-2.55	-3.21	-5.73 \pm 0.25	-4.86	-6.14	-31.5 \pm 1.1	-27.9	-33.8	14.3 \pm 1.3	16.1	9.3	27 \pm 9	47	3
Polychrome Hall	D4	22/03/ 17 to 12/08/ 2020	Droplets on the cave ceiling	V	-2.98 \pm 0.10	-2.78	-3.14	-5.69 \pm 0.19	-5.27	-5.98	-31.5 \pm 0.74	-30.4	-32.7	13.9 \pm 1.1	16.0	11.7	25 \pm 11	45	2
				S	-2.98 \pm 0.12	-2.67	-3.20	-5.68 \pm 0.23	-5.07	-6.14	-31.3 \pm 1.1	-28.5	-33.3	14.1 \pm 1.1	15.8	12.9	23 \pm 9	47	9
Polychrome Hall	D5	22/03/ 17 to 12/08/ 2020	Droplets on the cave ceiling	V	-3.01 \pm 0.12	-2.72	-3.28	-5.75 \pm 0.22	-5.23	-6.27	-31.7 \pm 1.0	-30.4	-34.7	14.3 \pm 1.1	16.5	11.2	27 \pm 10	45	10
				S	-2.99 \pm 0.11	-2.70	-3.20	-5.72 \pm 0.21	-5.14	-6.12	-31.2 \pm 1.0	-29.1	-33.4	14.5 \pm 1.0	17.3	11.1	29 \pm 10	59	11

445 (mod. 0560–4450) device with a precision of 5%. Relative humidity was measured using EE21 Series instrument, with an accuracy of $\pm 3\%$. Additionally, the automatic measurements performed by this device have being reproduced with grab sampling measurements carried out with an aspiration psicrometer Lambrecht kG Göttingen (model 761), with a resolution of 2%.

3.4. Stable isotope modeling

We modeled the stable isotope composition of condensation water at site C1 by using a two-endmember mixing model, in which the end-members are the water vapor generated within the cave (autochthonous water vapor hereafter) and the water vapor that enters the cave from outside during the cave ventilation period (allochthonous water vapor hereafter) (see Supplementary Table 2 for model parametrization and calculations). The CO_2 content in air is assumed to reflect the degree of atmosphere ventilation at site C1. This means that the maximum CO_2 content recorded at site C1 during the cave stagnation period (5525 ppm) represents 0% ventilation and the outside CO_2 content during the length of this study (470 ppm) represents 100% ventilation. We assume a linear relationship between the CO_2 content and the degree of cave ventilation at site C1. The isotopic composition of the two water vapor reservoirs can be approximated by assuming equilibrium conditions with environmental water (i.e. cave water and meteoric water). We use the mean isotopic composition of dripwater at site D1 to represent the isotopic composition of cave water ($\delta^{18}\text{O} = -5.9 \pm 0.1\text{‰}$, $\delta\text{D} = -32.8 \pm 0.4\text{‰}$, d-excess = $14.4 \pm 0.6\text{‰}$, ^{17}O -excess = 25 ± 7 per meg, during the length of this study). We use the long-term (i.e. 15 years mean) monthly values of $\delta^{18}\text{O}$ and δD in rainfall recorded by the meteorological station of Santander city, 15 km apart from Altamira Cave (data from the Global Network of Isotopes in Precipitation of the International Atomic Energy Agency, period 2000–2015; IAEA/WMO 2021) to calculate the isotopic composition of the external vapor. Because there is no $\delta^{17}\text{O}$ data, and therefore ^{17}O -excess cannot be calculated for precipitation near Altamira Cave, we assume that the ^{17}O -excess in rainfall correlates linearly with $\delta^{18}\text{O}$, as reported by Giménez et al. (2021) in the Pyrenees (northern Spain), ~ 220 km from Altamira Cave. These authors found that the $\delta^{18}\text{O}$ - ^{17}O -excess relationship has a slope of 3.8 per meg/‰ at this site. Then, we use this relationship to calculate the ^{17}O -excess of rainfall in the setting of Altamira Cave from the $\delta^{18}\text{O}$ of rainfall. Finally, the $\delta^{17}\text{O}$ is calculated from $\delta^{18}\text{O}$ and ^{17}O -excess.

The isotopic compositions of both, the autochthonous and the allochthonous water vapor are controlled by the in-cave and external temperatures, respectively. Equilibrium liquid–vapor fractionation factors for $\delta^{18}\text{O}$ ($\alpha^{18}\text{O}_{\text{eq}}$) and δD ($\alpha\text{D}_{\text{eq}}$) are calculated here as a function of temperature using the equations of Horita and Wesolowski (1994). The $\alpha^{17}\text{O}_{\text{eq}}$ is calculated as $\alpha^{17}\text{O}_{\text{eq}} = \alpha^{18}\text{O}_{\text{eq}}^\theta$, where θ is 0.529 ± 0.001 (Barkan and Luz, 2005).

4. Results

A total of 481 water samples from Altamira Cave were analyzed for triple oxygen and hydrogen isotopes. The summary of the $\delta^{17}\text{O}$, $\delta^{18}\text{O}$, δD values and derived parameters d-excess and ^{17}O -excess are given in Table 1 and Supplementary Table 3. The results are plotted with time in Fig. 2, along with the monitored climate parameters at site C1.

4.1. Condensation water from artificial supports

4.1.1. Outmost cave sector (site C1)

The $\delta^{18}\text{O}$ and δD values of condensation water from the outmost part of the cave correlate linearly ($\delta\text{D} = 6.9(\pm 0.2) \cdot \delta^{18}\text{O} + 9.1(\pm 1.1)$; $R^2 = 0.94$, p-value < 0.05). The slope of this relationship is similar to that of the Local Meteoric Water Line ($\delta\text{D} = 7.1(\pm 0.1) \cdot \delta^{18}\text{O} + 7.4(\pm 0.8)$; $R^2 = 0.93$, p-value < 0.05) in Santander city (15 km from Altamira Cave; IAEA/WMO, 2021), while the intercept is slightly higher than that of the

LMWL (Fig. 3A). The d-excess and ^{17}O -excess values of condensation water at site C1 are not correlated with $\delta^{18}\text{O}$ ($R^2 = 0.1$, p-value > 0.05) (Fig. 3). The $\delta^{18}\text{O}$ and δD values are higher during the cave ventilation period ($-4.6 \pm 0.7\text{‰}$ and $-22.4 \pm 4.6\text{‰}$, respectively), while during the rest of the year are lower and less variable ($-5.6 \pm 0.4\text{‰}$ and $-30.0 \pm 3.0\text{‰}$, respectively) (Figs. 2 and 4). The ^{17}O -excess values of condensation water at site C1 average 27 ± 11 per meg and ranges from 6 to 60 per meg. No systematic differences in this parameter have been between the ventilation period (27 ± 12 per meg) and the stagnation period (28 ± 9 per meg), probably due to insufficient analytical precision of the measurements (± 13 per meg, 1SD) (Fig. 4). The d-excess shows more variable values during the ventilation period ($14.3 \pm 2.1\text{‰}$) than during the cave stagnation period ($15.2 \pm 1.1\text{‰}$) (Figs. 2 and 4).

4.1.2. Inner cave sectors (sites C2 and C3)

The mean $\delta^{18}\text{O}$ and δD values of condensation water from the two sites investigated in the inner part of the cave (sites C2 and C3) are indistinguishable within errors (Table 1 and Supplementary Table 3). The $\delta^{18}\text{O}$ and δD values correlate linearly ($\delta\text{D} = 4.5(\pm 0.4) \cdot \delta^{18}\text{O} - 6.0(\pm 1.4)$; $R^2 = 0.76$, p-value < 0.05) (Fig. 3). The annual variability is < 0.3‰ (1SD) for $\delta^{18}\text{O}$ and < 1.6‰ for δD and does not correlate with seasonal changes in the monitored air parameters (temperature and CO_2 concentration) in the cave (Fig. 2). Indeed, no significant differences in all the measured isotopic parameters can be distinguished between the cave ventilation period and the stagnation period (Table 1 and Fig. 4). The correlation between d-excess and $\delta^{18}\text{O}$ is negative and statistically significant ($R^2 = 0.63$, p-value < 0.05) (Fig. 3). There are no significant correlations between ^{17}O -excess and $\delta^{18}\text{O}$ ($R^2 = 0.01$, p-value > 0.05) or between ^{17}O -excess and d-excess ($R^2 = 0.12$, p-value > 0.05) (Fig. 3).

4.2. Dripwater (site D1) and ceiling droplets (sites D2, D3, D4 and D5) in the Polychrome Hall

The $\delta^{18}\text{O}$ of the fast discharge point in the Polychrome Hall (site D1) ranged from -6.1 to -5.5‰ , with a mean of $-5.9 \pm 0.1\text{‰}$. The δD ranged from -33.7 to -31.6‰ , with a mean of $-32.8 \pm 0.4\text{‰}$. The d-excess varied between 11.9 and 15.0‰, with a mean of $14.4 \pm 0.6\text{‰}$, and the ^{17}O -excess varied between 11 and 39 per meg with a mean of 25 ± 7 per meg (Table 1). No clear systematic seasonal patterns have been observed in the isotopic parameters during the 4-years monitoring (Figs. 2 and 4).

The isotopic composition of the ceiling droplets with no apparent dripping in the Polychrome Hall (D2, D3, D4, D5) are statistically indistinguishable (Table 1 and Fig. 4). Taking altogether, the $\delta^{18}\text{O}$ of the ceiling droplets ranged from -6.3 to -4.7‰ , with a mean of $-5.7 \pm 0.2\text{‰}$. The δD ranged from -34.7 to -27.9‰ , with a mean of $-31.4 \pm 1\text{‰}$. The d-excess varied between 8.0 and 17.3‰, with a mean of $14.2 \pm 1.3\text{‰}$, and the ^{17}O -excess did between 2 and 61 per meg with a mean of 26 ± 10 per meg. No systematic seasonal patterns have been observed in the isotopic parameters during the 4-years monitoring (Table 1 and Fig. 4). The $\delta^{18}\text{O}$ and δD in the four points showed similar patterns with time ($R^2 > 0.4$ and p-value < 0.05).

4.3. Microclimate parameters

The air temperature in the Entrance Hall (site C1 in Fig. 1) ranged from 13.1 to 15.7°C , with a mean value of $14.5 \pm 0.6^\circ\text{C}$. Maximum temperatures were recorded in September and minimum temperatures correspond to March. In the Polychrome Hall (sites D1, D2, D3, D4 and D5 in Fig. 1) air temperature oscillated between 13.2 and 15.0°C and the mean value was $14.2 \pm 0.5^\circ\text{C}$. Maximum temperatures were recorded in December, while minimum temperatures correspond to June. The temperature in the innermost sector of the cave (locations C2 and C3 in Fig. 1) varied between 12.5 and 14.5°C , with a mean of $13.7 \pm 0.2^\circ\text{C}$. Maximum temperatures were recorded between October and February and minimum temperatures corresponded to June and July.




Fig. 3. Cross-plot of isotopic parameters (A. δD vs. $\delta^{18}O$; B. d-excess vs. $\delta^{18}O$; C. ^{17}O -excess vs. $\delta^{18}O$; D. ^{17}O -excess vs. d-excess) in condensation waters (sites C1, C2 and C3), dripwater (D1) and ceiling droplets with no apparent dripping in the ceiling of the *Polychrome Hall* (D2, D3, D4 and D5) of Altamira Cave. The local meteoric water line (LMWL; IAEA/WMO, 2021) is given for comparison in panel A. The errors for $\delta^{18}O$ and δD are smaller than the symbols. The mean analytical errors for d-excess (0.2‰) and ^{17}O -excess (13 per meg) are represented in panels B, C and D.

The temperature outside measured during the visits to the cave (between 10 a.m. and 12 p.m.) ranged from 2.9 to 25.1 °C, with a mean of 15.1 ± 4.7 °C. Maximum temperatures outside were recorded in August, while minimum temperatures correspond to January and February. August and July are the driest month of the series, while the autumn and spring months concentrate most of the annual precipitation (Fig. 2).

The mean CO_2 in the *Polychrome Hall* was 2580 ± 1423 ppm, with lower values around 1000 ppm from June to October (“ventilation period”) and higher values around 3000 ppm were recorded between November and May (“stagnation period”). At site C1, the CO_2 content varied between 724 and 5525 ppm. The change from lower to higher CO_2 conditions (and vice versa) generally takes place in around 2 weeks, mainly depending on the external meteorology.

Relative humidity (RH) in the *Polychrome Hall* was $96.8 \pm 1.9\%$ on average and ranged from 92% to 100%. In the outmost part of the cave (site C1), RH ranged from 88% to 100% and averaged $96.5 \pm 2.3\%$ (Fig. 3E). Generally, the lowest RH values correspond to the cave ventilation period and values closer to 100% are characteristic of the stagnation period. The RH in the inner cave sector is slightly higher and closer to saturation (>97%), with no significant seasonal changes (Gázquez et al., 2016; Sainz et al., 2018).

4.4. Results of isotope modeling

The modeled $\delta^{18}O$ and δD values of condensation water for the sectors near the cave entrance (site C1) show a seasonal pattern, with higher values (up to -2.9% and -8.4% , respectively) during the cave ventilation period and lower values during the stagnation period (down to -6.9% and -43.7% , respectively) (Fig. 1). The modeled d-excess values of condensation water ranges from 10.1 to 16.2‰, with the lowest values corresponding to the ventilation period and higher values are obtained for the stagnation period. We found that the expected range of ^{17}O -excess in condensation water is only of 9 per meg. Note that this variability is smaller than the analytical precision of our ^{17}O -excess measurements (~ 13 per meg). The highest ^{17}O -excess values occur during the cave stagnation period (maximum value of 28 per meg) and lower values are observed during the cave ventilation period (minimum of 19 per meg).

We tested the sensibility of the isotopic composition of condensation water to the different parameters consider by our model (external air temperature and cave atmosphere temperature, the isotopic composition of cave dripwater and meteoric water and the degree of cave atmosphere ventilation). As expected, we found that during the cave stagnation period the isotopic composition of condensation water at site C1 is insensitive to changes in the outside air temperature and variations




Fig. 4. Box and whisker plots of $\delta^{18}\text{O}$ (A), δD (B), d-excess (C) and ^{17}O -excess (D) of condensation water in the outmost cave sector (sites C1) and in the inner sectors (sites C2 and C3), as well as dripwater (DW, site D1) and ceiling droplets with no apparent dripping (sites D2, D3, D4 and D5) in the *Polychrome Hall* of Altamira Cave. Results of the cave ventilation period (V; June to October) and the stagnation period (S; November to May) are displayed for comparison.

in the isotopic composition of meteoric water (Supplementary Figs. 1 and 3). Also, changes in the cave atmosphere temperature during the stagnation period have an insignificant impact on the isotopic composition of condensation water (Supplementary Fig. 2), given the small variations in the cave atmosphere temperature ($<5^\circ\text{C}$). In contrast, the isotopic composition of condensation water at site C1 during the cave ventilation period is relatively sensitive to the changes in the external air temperature (e.g. 1‰ $\delta^{18}\text{O}$ increase for each increment of 10°C) (Supplementary Fig. 1). We found that the $\delta^{18}\text{O}$ and δD values of condensation water increase linearly with the $\delta^{18}\text{O}$ and δD values of the external meteoric water (Supplementary Fig. 3), while changes in the cave temperature atmosphere have little impact on the $\delta^{18}\text{O}$ and δD values of condensation water (e.g. 0.2‰ $\delta^{18}\text{O}$ increase for an increment of 2°C) (Supplementary Fig. 2). Likewise, the secondary parameters d-excess and ^{17}O -excess are insensitive to change in the external parameters during the cave stagnation period (Supplementary Figs. 1 to 3). In contrast, during the ventilation period, the d-excess of condensation water in the outmost cave sectors is significantly impacted by changes in the external air temperature (i.e. 3.7‰ d-excess increase for each increment of 10°C) and by changes in the d-excess of meteoric water. The modeled ^{17}O -excess values of condensation water at site C1 during the ventilation period are barely affected by changes in the internal and external air temperatures (i.e. 1 per meg increase for each increment of 10°C).

5. Discussion

The difference in isotopic composition of two water reservoirs (e.g. liquid and vapor) in equilibrium during phase change depends mostly on temperature that controls the equilibrium isotope fractionation factors between the two phases (e.g. $\alpha^{18}\text{O}_{\text{water-vapor}}$ and $\alpha\text{D}_{\text{water-vapor}}$) (Craig et al., 1963; Horita and Wesolowski, 1994). Additionally, when the evaporation–condensation process occurs under $\text{RH} < 100\%$, kinetic isotope fractionation can take place, which magnitude depends mostly on relative humidity and the degree of turbulence on the liquid–vapor boundary layer (i.e. wind) (Cappa et al., 2003; Landais et al., 2006; Barkan and Luz, 2007; Luz et al., 2009). In caves, relative humidity is

close to 100% (e.g. Fig. 2) and wind speed is generally insignificant compared to outside (Gázquez et al., 2016); thus, equilibrium isotope process is expected to prevail over kinetic isotope fractionation during phase changes. Consequently, for the particular case of condensation in caves, the most important parameters controlling the isotopic composition of condensation water are temperature and the isotopic composition of water vapor.

Our monitoring data show that Altamira Cave directly, and rapidly, exchanges gases with the external atmosphere during the cave ventilation period (June to October), when intakes of water vapor from outside are expected. Note that the seasonal environmental changes outside are considerably larger than in the cave. This includes larger temperature changes and varying isotopic compositions of water vapor. Indeed, the isotopic composition of summer rainfall in this region is more enriched (e.g. $\delta^{18}\text{O} \sim -3.5\text{‰}$ in June) than in winter (e.g. $\delta^{18}\text{O} \sim -6.7\text{‰}$ in January) (IAEA/WMO, 2021) and external water vapor is assumed here to be in equilibrium with environmental liquid water outside. According to this, the isotopic composition of atmospheric vapor outside the cave during summer is expected to increase compared to winter conditions. In contrast, the isotopic composition of the autochthonous water vapor generated within the cave is expected to be less variable than outside because of cave temperature is nearly constant and the vapor is in equilibrium with the cave dripwater, which has almost constant isotopic values (Table 1 and Fig. 2). We evaluate the impact of allochthonous water vapor entering the cave during the ventilation period on the isotopic composition of condensation water at site C1 (*Entrance Hall*) by using a two-endmember mixing model (Fig. 2 and Supplementary Table 2).

The magnitude and variability of the modeled isotopic values for site C1 agree with our observations of $\delta^{18}\text{O}$ and δD of condensation water at site C1 (Fig. 2). Our model demonstrates that the seasonal cycles recorded by the isotopic composition of condensation water in the outmost part of the cave are a consequence of advection of allochthonous water vapor from the external atmosphere. The “summer condensation water” that accumulates during the ventilation period near the *Entrance Hall* is isotopically enriched (i.e. higher $\delta^{18}\text{O}$ and δD values) compared to the “winter condensation water” that derives from

recycling of autochthonous water vapor generated in the cave water during the stagnation period (Figs. 3 and 5). The relatively constant $\delta^{18}\text{O}$ and δD values of autochthonous water vapor (ranges of 0.3‰ and 2.7‰, respectively) is a consequence of the small cave temperature variability and the almost constant isotopic values of seepage water through the seasonal cycle.

Condensation from allochthonous water vapor in caves seems to be analogue to recharge by “occult rain” detected in arid regions that occurs because of re-evaporation of local water and subsequent condensation of vapor on vegetation and soils (Aravena et al., 1989; Kaseke et al., 2017). If such vapor is re-condensed in any significant quantity before mixing with the larger tropospheric reservoir, the isotopic composition of the resulting condensation water will fall slightly above the LMWL (i.e. higher d-excess values), along a condensation line with a slope similar to the LMWL (Ingraham and Matthews, 1988, 1990; Clark and Fritz, 1997). Indeed, we found that y-intercept of the $\delta^{18}\text{O}$ - δD correlation line of condensation water in the outmost part of the cave (site C1) is slightly higher ($9.1 \pm 1.1\text{‰}$) than that of the Local Meteoric Water Line (LMWL) ($7.4 \pm 0.8\text{‰}$), although their slopes are indistinguishable within errors (Fig. 3). Similarly, an offset between d-excess of condensation water and the LMWL has been observed in other caves (Gázquez et al., 2017b; Liñán et al., 2021).

Evaporation of condensation water can explain some of the d-excess outliers observed in the *Entrance Hall* that were recorded mostly during the cave ventilation period (e.g. three d-excess values ranging from 7 to

10.5‰ in summer 2017, 2018 and 2019; Fig. 4). These d-excess values are slightly lower than predicted by our model (Fig. 2C). Note that evaporated water normally show lower d-excess values (that correlates with $\delta^{18}\text{O}$ and δD) compared to non-evaporated waters (e.g. Luz et al., 2009; Gázquez et al., 2018; Voigt et al., 2021). However, the correlation between $\delta^{18}\text{O}$ and d-excess of condensation water in Altamira Cave is weak ($R^2 = 0.28$), the d-excess of condensation water remains generally high during the ventilation period ($\sim 15\text{‰}$ and up to 17.3‰) and can occasionally show lower values during the rest of the year. In consequence, the seasonal variability of condensation water at site C1 cannot be attributed to the effect of evaporation, but to the varying contributions of autochthonous and allochthonous water vapor.

The modeled ^{17}O -excess values of condensation water at site C1 vary from 19 per meg during the cave stagnation period to 28 per meg during the ventilation period (Fig. 2). No statistically significant seasonal trends in ^{17}O -excess of our analytical data have been observed in any of the sites investigated in this study, either in condensation water nor in dripwater. Recent studies showed that the $\delta^{18}\text{O}$ - ^{17}O -excess relationship in rainwater in northern Spain (Villanua village, ~ 220 km from Altamira Cave) has a slope of 3.8 per meg/‰. Considering that the $\delta^{18}\text{O}$ of rainwater in Santander city varies seasonally by $\sim 3\text{‰}$, a range of ^{17}O -excess in rainwater of 12 per meg would be expected. The expected ^{17}O -excess variability in dripwater of Altamira Cave is even smaller (2 per meg), considering the small $\delta^{18}\text{O}$ variability observed during the length of this study (0.5‰). Regarding that the mean reproducibility of our ^{17}O -excess measurements is 13 per meg (1SD), we conclude that the current analytical precision of the CRDS analyzers is insufficient to investigate the small ^{17}O -excess variability expected for condensation water and dripwater in caves.

As for condensation water in the inner cave sectors, the triple oxygen and hydrogen isotopic values in dripwater and its temporal variability (site D1) are indistinguishable from those of condensation water collected from the inner cave sectors (sites C2 and C3). This suggests that water condensation from autochthonous water vapor in Altamira Cave cannot be differentiated from seepage water by using oxygen and hydrogen isotopes. Also, the isotopic composition of the ceiling droplets with no apparent dripping (sites D2, D3, D4 and D5) of the *Polychrome Hall* and the rest of condensation waters from the inner cave sectors (sites C2 and C3) show indistinguishable values within errors. No clear differences between the ventilation and the stagnation periods has been identified. Thus, it is possible that the ceiling droplets with of the *Polychrome Hall* are related to water condensation; however, contributions of meteoric water infiltration and extremely slow drainage of the epikarst over this chamber cannot be ruled out from our data.

We found that the small seasonal changes observed in the isotopic composition of dripwater and condensation water in sectors relatively distant from the cave entrance do not seem to be controlled by ventilation/stagnation regimen of the cave, since no correlation with the CO_2 content or with in-cave air temperature has been observed (Fig. 4). This indicates that the main source of vapor for condensation in these sites is autochthonous water vapor from within the cave, with insignificant contributions of allochthonous air masses, even during the ventilation period. This is probably related to the fact that allochthonous water vapor condenses on cave surfaces closer to the entrance and, likely on the artificial door that separates the *Entrance Hall* and the rest of the cave, which is permanently closed. In consequence, the allochthonous water vapor does not reach the inner cave sector even during the ventilation periods and does not impact on the isotopic composition of condensation water in these areas.

6. Conclusions

We investigate the spatiotemporal variability of $\delta^{17}\text{O}$, $\delta^{18}\text{O}$, δD and derived ^{17}O -excess and d-excess values of condensation water in Altamira Cave. Significant changes in some of these parameters have been observed in the outmost cave sectors, while in the inner/deeper cave




Fig. 5. Conceptual diagram of condensation water systematic in Altamira Cave, during the cave ventilation period (A) and the cave stagnation period (B). The inner cave sectors are less affected by water vapor transported by air masses from outside during the entire year, while the main source of vapor for condensation in these areas is autochthonous water vapor generated from dripwater. The outmost areas receive allochthonous water vapor from outside only during the ventilation period, while the autochthonous vapor is the main source of moisture for condensation in the entire cave during the stagnation period. The metal door that separates the *Entrance Hall* from the rest of the cave has been represented.

parts the isotopic composition of condensation water is less variable though the seasonal cycles. In particular, we found that the $\delta^{18}\text{O}$ and δD of condensation water in the cave sectors closer to the entrance during the ventilation period (June to October) are higher than during the stagnation period (November to May). We evaluate the role of cave ventilation pattern in the condensation process in this cave sector by using an isotope mixing model. We found that the isotopic composition of condensation water near the cave entrance during the ventilation period agrees with the expected values of water condensed from allochthonous water vapor, while the prevailing mechanism during the stagnation period is condensation from autochthonous water vapor generated within the cave.

Importantly, no significant differences have been found between the isotopic composition of condensation water in the inner cave sectors and dripwater. This suggests that the water condensation process in areas more distant from the cave entrance is sourced by autochthonous water vapor. Given that the isotopic composition of condensation water though the seasonal cycle is indistinguishable from that of seepage water in the cave, we conclude that oxygen and hydrogen isotopes are not useful in identifying the origin (condensation vs. seepage) of the droplets with no apparent dripping that cover the ceiling of the *Polychrome Hall*. Also, we found that the analytical precision of the current CRDS isotope analyzers to measure triple oxygen isotope anomalies in waters is not enough to resolve the small natural seasonal variations expected for cave waters (tens per meg).

The varying contributions of the allochthonous and autochthonous water vapor sources for condensation in the cave can be attributed the natural cave ventilation, as well as to the distance of the different chambers to the cave entrance and probably to the role in the cave microclimate of the artificial metal door that separates the *Entrance Hall* and the rest of the cave, as observed in previous studies (Gázquez et al., 2016; Sainz et al., 2017). It is possible that a decrease in the airflow speed during the ventilation period in this cave sector favors that water vapor from outside condenses mostly in surfaces near the artificial door. Thus, this artificial element seems to be critical to prevent the *Polychrome Hall* and the rest of the cave from receiving additional water vapor from outside during the ventilation period. Our results are important for the management of the *Polychrome Hall*, since condensation water can contribute to the deterioration of its exceptional rock art by washing out the ochre pigments. Additional efforts should be made to quantify and monitor condensation processes in Altamira Cave in forthcoming years.

Author Contribution Statement

FG conducted the experimental design of the study and wrote the manuscript. LQ, DR, IF, SC and CS performed the water sampling, the cave monitoring and collaborated in the preparation of the manuscript.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors are grateful to the Altamira Cave Research Centre and Museum staff for their help and support during the sampling surveys in the cave. The image of the ceiling of the *Polychrome Hall* was provided by the Altamira Cave Research Center. This work was supported by the Ministry of Education, Culture and Sport of Spain (MECD) [grant number J180052] under the Project “Servicio de control y seguimiento de las condiciones ambientales y del biodeterioro de la Cueva de Altamira”, and also with the support of the Cantabrian International Institute for Prehistoric Research (IIIPC) under the project entitled “Control y

seguimiento de las condiciones ambientales, del agua de infiltración y de las emisiones de gas radon, de la cueva de Altamira”. The authors thanks Professor David A. Hodell for the isotopic analysis of 15 water samples. Dr. Fernando Gázquez was financially supported by the HIPATIA research program of the University of Almería and by a Ramón y Cajal Fellowship (RYC2020-029811-I) of the Spanish Government (Ministerio de Ciencia e Innovación). The authors are grateful to four anonymous reviews, who contributed to improve the original manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jhydrol.2021.127416>.

References

- Aravena, R., Suzuki, O., Polastri, A., 1989. Coastal fogs and their relation to groundwater. *Chem. Geol.* 79, 83–91.
- Aron, P.G., Levin, N.E., Beverly, E.J., Huth, T.E., Passey, B.H., Pelletier, E.M., Poulsen, C. J., Winkelstern, I., Yarian, D., 2021. Triple oxygen isotopes in the water cycle. *Chem. Geol.* 565 (1), 120026.
- Auler, A.S., Smart, P.L., 2004. Rates of condensation corrosion in speleothems of semi-arid northeastern Brazil. *Speleogenesis Evol. Karst Aquifers* 2, 2.
- Barkan, E., Luz, B., 2005. High precision measurements of $^{17}\text{O}/^{16}\text{O}$ and $^{18}\text{O}/^{16}\text{O}$ ratios in H_2O . *Rapid Commun. Mass Spectrom.* 19 (24), 3737–3742.
- Barkan, E., Luz, B., 2007. Diffusivity fractionations of $\text{H}_2^{16}\text{O}/\text{H}_2^{17}\text{O}$ and $\text{H}_2^{16}\text{O}/\text{H}_2^{18}\text{O}$ in air and their implications for isotope hydrology. *Rapid Commun. Mass Spectrom.* 21 (18), 2999–3005.
- Bowen, G.J., Cai, Z., Fiorella, R.P., Putman, A.L., 2019. Isotopes in the Water Cycle: Regional- to Global-Scale Patterns and Applications. *Ann. Rev. Earth Plant. Sci.* 47 (1), 453–479.
- Calaforra, J.M., 1996. Contribución al conocimiento de la karstología de yesos (PhD Thesis). University of Granada, Spain, p. (350 pp.).
- Calaforra, J.M., Dell’Aglia, A., Forti, P., 1993. Preliminary data on the chemical erosion in gypsum karst. The Sorbas region (Spain). In: *Proceedings of the 11th International Congress of Speleology* (Beijing), pp. 97–99.
- Cappa, C.D., Hendricks, M.B., DePaolo, D.J., Cohen, R.C., 2003. Isotopic fractionation of water during evaporation. *J. Geophys. Res.* 108 (D16), 4525.
- Clark, I.D., Fritz, P., 1997. *Environmental Isotopes in Hydrogeology*. Lewis Publishers, USA.
- Craig, H., Gordon, L.I., Horibe, Y., 1963. Isotopic exchange effects in the evaporation of water: 1. Low-temperature experimental results. *J. Geophys. Res.* 68 (17), 5079–5087.
- Cuezva, S., Fernandez-Cortes, A., Benavente, D., Serrano-Ortiz, P., Kowalski, A.S., Sanchez-Moral, S., 2011. Short-term $\text{CO}_2(\text{g})$ exchange between a shallow karstic cavity and the external atmosphere during summer: Role of the surface soil layer. *Atmos. Environ.* 45 (7), 1418–1427.
- Cuezva, S., Sanchez-Moral, S., Saiz-Jimenez, C., Cañaveras, J.C., 2009. Microbial communities and associated mineral fabrics in Altamira Cave, Spain. *Int. J. Speleol.* 38 (1), 83–92.
- De Freitas, C.R., Schmokal, A.A., 2003. Condensation as a microclimate process: measurement, numerical simulation and prediction in the Glowworm tourist cave, New Zealand. *Int. J. Climatol.* 23, 557–575.
- De Freitas, C.R., Schmokal, A., 2006. Studies of corrosion/condensation process in the Glowworm Cave, New Zealand. *Int. J. Speleol.* 35, 75–81.
- Dreybrodt, W., Gabrovšek, F., Perne, M., 2005. Condensation corrosion: a theoretical approach. *Acta Carsol.* 34, 317–348.
- Dublyansky, V.N., Dublyansky, Y.V., 1998. The problem of condensation in karst studies. *J. Caves Karst Stud.* 60 (1), 3–17.
- Dublyansky, V.N., Dublyansky, Y.V., 2000. The role of condensation in karst hydrogeology and speleogenesis. In: Klimchouk, A.B., Ford, D.C., Palmer, A.N., Dreybrodt, W. (Eds.), *Speleogenesis, Evolution of Karst Aquifers*. National Speleological Society, Huntsville, pp. 100–111.
- Elez, J., Cuezva, S., Fernández-Cortés, A., García-Antón, E., Benavente, D., Cañaveras, J. C., Sánchez-Moral, S., 2013. A GIS-based methodology to quantitatively define an Adjacent Protected Area in a shallow karst cavity: case of Altamira cave. *J. Environ. Manage.* 118, 122–124.
- Fernández-Cortés, A., Calaforra, J.M., García-Guinea, J., 2006. The Pulpí gigantic geode (Almería, Spain): geology, metal pollution, microclimatology and conservation. *Environ. Geol.* 50 (5), 707–716.
- Gabrovšek, F., Dreybrodt, W., Perne, M., 2010. Physics of condensation corrosion in caves. In: Andreo, B., Carrasco, F., Durán, J.J., LaMoreaux, J.W. (Eds.), *Advances in Research in Karst Media, Environmental Earth Sciences*, pp. 491–499.
- García-Antón, E., Cuezva, S., Jurado, V., Porca, E., Miller, A.Z., Fernández-Cortés, A., Saiz-Jiménez, C., Sánchez-Moral, S., 2013. Combining stable isotope ($\delta^{13}\text{C}$) of trace gases and aerobiological data to monitor the entry and dispersion of microorganisms in caves. *Environ. Sci. Poll. Res.* 21, 473–484.
- García-Antón, E., Cuezva, S., Fernandez-Cortes, A., Benavente, D., Sanchez-Moral, S., 2014. Main drivers of diffusive and advective processes of CO_2 -gas exchange between a shallow vadose zone and the atmosphere. *Int. J. Greenh. Gas. Con.* 21, 113–129.

- Gázquez, F., Calaforra, J.-M., Forti, P., De Waele, J.o., Sanna, L., 2015a. The role of condensation in the evolution of dissolutional forms in gypsum caves: study case in the karst of Sorbas (SE Spain). *Geomorphology* 229, 100–111.
- Gázquez, F., Mather, I., Rolfe, J., Evans, N.P., Herwartz, D., Staubwasser, M., Hodell, D. A., 2015b. Simultaneous analysis of $^{17}\text{O}/^{16}\text{O}$, $^{18}\text{O}/^{16}\text{O}$ and $^2\text{H}/^1\text{H}$ of gypsum hydration water by cavity ringdown laser spectroscopy. *Rapid Commun. Mass Spectrom.* 29 (21), 1997–2006.
- Gázquez, F., Quindós-Ponceta, L., Sainz, C., Fernández, A., Fuentes, I., Celaya, S., 2016. Spatiotemporal distribution of $^{13}\text{C}\text{-CO}_2$ in a shallow cave and its potential use as indicator of anthropic pressure. *J. Environ. Manage.* 180, 421–432.
- Gázquez, F., Rull, F., Sanz-Arranz, A., Medina, J., Calaforra, J.M., de las Heras, C., Lasheras, J.A., 2017a. In situ Raman characterization of minerals and degradation processes in a variety of cultural and geological heritage sites. *Spectrochim. Acta Part B* 172, 48–57.
- Gázquez, F., Calaforra, J.M., Evans, N.P., Hodell, D.A., 2017b. Using stable isotopes ($\delta^{17}\text{O}$, $\delta^{18}\text{O}$ and δD) of gypsum hydration water to ascertain the role of water condensation in the formation of subaerial gypsum speleothems. *Chem. Geol.* 452, 34–46.
- Gázquez, F., Morellón, M., Bauska, T., Herwartz, D., Surma, J., Moreno, A., Staubwasser, M., Valero-Garcés, B., Delgado-Huertas, A., Hodell, D.A., 2018. Triple oxygen and hydrogen isotopes of gypsum hydration water for quantitative paleo-humidity reconstruction. *Earth Planet. Sci. Lett.* 481, 177–188.
- Giménez, R., Bartolomé, M., Gázquez, F., Iglesias, M., Moreno, A., 2021. Underlying climate controls in triple oxygen (^{16}O , ^{17}O , ^{18}O) and hydrogen (^1H , ^2H) isotopes composition of rainfall (Central Pyrenees). *Front. Earth Sci.* 9, 633698.
- Horita, J., Wesolowski, D.J., 1994. Liquid–vapor fractionation of oxygen and hydrogen isotopes of water from the freezing to the critical temperature. *Geochim. Cosmochim. Acta* 58 (16), 3425–3437.
- Hoyos, M., Bustillo, A., García, A., Martín, C., Ortiz, R., Suazo, C., 1981. Características geológico-kársticas de la cueva de Altamira (Santillana del Mar, Santander). Informe Ministerio de Cultura, Madrid, p. 81.
- IAEA/WMO (2021). Global Network of Isotopes in Precipitation. The GNIP Database. Accessible at: <http://www.iaea.org/water>.
- Ingraham, N.L., Matthews, R.A., 1988. Fog drip as a source of groundwater recharge in northern Kenya. *Water Resour. Res.* 24 (8), 1406–1410.
- Ingraham, N.L., Matthews, R.A., 1990. A stable isotopic study of fog: the Point Reyes Peninsula, California, U.S.A. *Chem. Geol.* 80 (4), 281–290.
- Kaseke, K.F., Wang, L., Seely, M.K., 2017. Nonrainfall water origins and formation mechanisms. *Sci. Adv.* 3, e1603131.
- Klimchouk, A.B., 1996. Speleogenesis in gypsum. *Int. J. Speleol.* 25 (3–4), 61–82.
- Klimchouk, A.B., Aksem, S.D., 2002. Gypsum karst in the western Ukraine: Hydrochemistry and solution rates. *Carbonate Evaporite* 17, 142–153.
- Landais, A., Barkan, E., Yakir, D., Luz, B., 2006. The triple isotopic composition of oxygen in leaf water. *Geochim. Cosmochim. Acta* 70 (16), 4105–4115.
- Liñán, C., Benavente, J., del Rosal, Y., Vadillo, I., Ojeda, L., Carrasco, F., 2021. Condensation water in heritage touristic caves: isotopic and hydrochemical data and a new approach for its quantification through image analysis. *Hydrol. Process.* 35 (5) <https://doi.org/10.1002/hyp.v35.510.1002/hyp.14083>.
- Luz, B., Barkan, E., 2010. Variations of $^{17}\text{O}/^{16}\text{O}$ and $^{18}\text{O}/^{16}\text{O}$ in meteoric waters. *Geochim. Cosmochim. Acta* 74 (22), 6276–6286.
- Luz, B., Barkan, E., Yam, R., Shemesh, A., 2009. Fractionation of oxygen and hydrogen isotopes in evaporating water. *Geochim. Cosmochim. Acta* 73 (22), 6697–6703.
- Pike, A.W.G., Hoffmann, D.L., García-Ideiz, M., Pettitt, P.B., Alcolea, J., Balbín, R.D., González-Saiz, C., De las Heras, C., Lasheras, J.A., Montes, R., Zilhao, J., 2012. U-Series Dating of Paleolithic Art in 11 Caves in Spain. *Science* 336 (6087), 1409–1413.
- Uechi, Y., Uemura, R., 2019. Dominant influence of the humidity in the moisture source region on the ^{17}O -excess in precipitation on a subtropical island. *Earth Planet. Sci. Lett.* 513, 20–28.
- Quindós, L.S., Bonet, A., Díaz-Caneja, N., Fernandez, P.L., Gutierrez, I., Solana, J.R., Soto, J., Villar, E., 1987. Study of the environmental variables affecting the natural preservation of the Altamira Cave paintings located at Santillana del Mar, Spain. *Atmos. Environ.* 21 (3), 551–560.
- Sainz, C., Rábago, D., Celaya, S., Fernández, E., Quindós, J., Quindós, L., Fernández, A., Fuente, I., Artech, J.L., Quindós, L.S., 2018. Continuous monitoring of radon gas as a tool to understand air dynamics in the cave of Altamira (Cantabria, Spain). *Sci. Total Environ.* 624, 416–423.
- Sarbu, S.M., Lascu, C., 1997. Condensation Corrosion in Movile Cave. Romania. *J. Caves Karst Stud.* 59, 99–102.
- Schoenemann, S.W., Schauer, A.J., Steig, E.J., 2013. Measurement of SLAP2 and GISP $\delta^{17}\text{O}$ and proposed VSMOW-SLAP normalization for $\delta^{17}\text{O}$ and ^{17}O excess. *Rapid Commun. Mass Spectrom.* 27 (5), 582–590.
- Steig, E.J., Gkinis, V., Schauer, A.J., Schoenemann, S.W., Samek, K., Hoffnagle, J., Dennis, K.J., Tan, S.M., 2014. Calibrated high-precision ^{17}O -excess measurements using cavity ring-down spectroscopy with laser-current-tuned cavity resonance. *Atmos. Meas. Tech.* 7, 2421–2435.
- Surma, J., Assonov, S., Staubwasser, M., 2021. Triple Oxygen Isotope Systematics in the Hydrologic Cycle. *Rev. Mineral. Geochem.* 86 (1), 401–428.
- Tarhule-Lips, R.F.A., Ford, D.C., 1998. Condensation corrosion in caves on Cayman Brac and Isla de Mona. *J. Caves Karst Stud.* 60, 84–95.
- Tian, C., Wang, L., Kaseke, K.F., Bird, B.W., 2018. Stable isotope compositions (d^2H , d^{18}O and d^{17}O) of rainfall and snowfall in the central United States. *Sci. Rep.* 8, 6712.
- Voigt, C., Herwartz, D., Dorador, C., Staubwasser, M., 2021. Triple oxygen isotope systematics of evaporation and mixing processes in a dynamic desert lake system. *Hydrol. Earth Syst. Sci.* 25, 1211–1228.
- Wassenaar, L., Terzer-Wasmuth, S., Douence, C., 2021. Progress and challenges in dual- and triple-isotope ($\delta^{18}\text{O}$, $\delta^2\text{H}$, $\Delta^{17}\text{O}$) analyses of environmental waters: An international assessment of laboratory performance. *Rapid Commun. Mass Spectrom.* 35 (24), e9193.
- White, J.H., Domínguez-Villar, D., Hartland, A., 2021. Condensation corrosion alters the oxygen and carbon isotope ratios of speleothem and limestone surfaces. *Res. Geochem.* e100008.